

COLLEGE OF HEALTH SCIENCES SCHOOL OF MEDICINE DEPARTMENT OF ANAESTHESIA

ANTIMICROBIAL PROPHYLAXIS IN CAESAREAN SECTION IN MULAGO HOSPITAL: EFFECT OF ADMINISTRATION TIME ON INCIDENCE OF POST-OPERATIVE INFECTION: RANDOMISED CLINICAL TRIAL

INVESTIGATOR DR. LOMANGISI DIANA DLAMINI

(MBChB, Medunsa)

A DISSERTATION TO BE SUBMITTED IN PARTIAL FULFILLMENT OF THE REQUIREMENTS FOR THE AWARD OF THE DEGREE OF MASTER OF MEDICINE (ANAESTHESIOLOGY AND CRITICAL CARE) OF MAKERERE UNIVERSITY

JUNE 2014

DECLARATION

| declare that this study is original and has not been published and/or submitted for any other degree award to any university before. |
|--|
| Signed Date |
| Dr Lomangisi Diana Dlamini |
| This dissertation has been submitted for examination with approval of the following supervisors: |
| Dr Arthur Kwizera |
| MBChB, MMed (Anaesthesiology) |
| Signed Date |
| Dr Musa Sekikubo |
| MBChB, MMed (OBS & GYN) |
| Signed Date |

DEDICATION

This book is dedicated to all the special women in my life past, present and future generation for a better health for all.

ACKNOWLEDGEMENT

My family - husband Mr Deogratius Sserumaga and daughter Charlene thank you for the support you gave me throughout the study.

My supervisors Dr Arthur Kwizera and Dr Musa Sekikubo thank you for the support and advice on this study.

My faithful and hard-working research assistants Jjuuko David and Kaweesa Isaac you worked tirelessly to give complete results your work is appreciated.

My statistician Dr Humphrey Wanzira thanks you for the good work.

To the Labour suite theatre team thank you for the support.

Above all praise be to God for the gift of life, strength and wisdom He gave me throughout the study.

Table of Contents

| DECLARATION | ii |
|-------------------------|------|
| DEDICATION | iii |
| ACKNOWLEDGEMENT | iv |
| List of tables | vii |
| List of figures | viii |
| List of appendices | ix |
| Operational Definitions | X |
| ABSTRACT | Xi |
| CHAPTER ONE | 1 |
| BACKGROUND | 1 |
| Research Problem | 2 |
| Research Question | 3 |
| Objectives | 3 |
| General Objective | 3 |
| Specific Objective | 3 |
| Hypotheses | 4 |
| Justification | 4 |
| CHAPTER TWO | 5 |
| LITERATURE REVIEW | 5 |
| CHAPTER THREE | 9 |
| METHODOLOGY | 9 |
| Study design | 9 |
| Study setting | 9 |
| Study population | 9 |
| Sampling method | 10 |
| Sample size | 11 |
| Trial registration | 12 |
| Data Management | 12 |
| Data collection | 12 |
| Data entry | 13 |
| Data analysis | |
| Ethical Consideration | |

| CHAPTER FOUR | 15 |
|---|----|
| RESULTS | 15 |
| CHAPTER FIVE | 20 |
| DISCUSSION | 20 |
| Study Limitations | 22 |
| CHAPTER SIX | 23 |
| CONCLUSION | 23 |
| References | 24 |
| APPENDICES | 27 |
| APPENDIX 1: Criteria for Defining a Surgical Site Infection (SSI) | 27 |
| APPENDIX 2: Consent Form (ENGLISH) | 29 |
| APPENDIX 3: Consent Form (LUGANDA) | 32 |
| APPENDIX 4: Data Collecting Tool | 35 |
| APPENDIX 5: Approval Letter | 37 |

List of tables

| Table 1. Distribution and characteristics of patients in the two groups | 16 |
|--|--------|
| Table 2. Mean time between incision and prophylaxis and average length of operation in | ı both |
| groups | 18 |
| Table 3. Outcome showing incidence of infection on both groups and neonatal outcomes | s18 |
| Table 4. Measure of association of incidence of infection with significant general | |
| characteristics | 19 |
| Table 5. Criteria for Defining a Surgical Site Infection (SSI)* | 27 |

List of figures

| Figure 1. Patient flow diagram | 15 |
|--|----|
| Figure 2. Distribution of indication for caesarean section in the two groups | 17 |
| | |
| Figure 3. Distribution of parity in the two groups | 17 |

List of appendices

Appendix 1: Table for defining surgical site infection

Appendix 2: Consent form English

Appendix 3: Consent form Luganda

Appendix 4: Data collecting tool

Appendix 5: Approval letter

Operational Definitions

WHO: World Health Organisation

Prophylactic antibiotics: antimicrobials given before contamination or infection to prevent

post-operative infection

Infection: Invasion by and multiplication of pathogenic microorganisms in a bodily part or

tissue, which may produce subsequent tissue injury and progress to overt disease through a

variety of cellular or toxic mechanisms.

Early post-operative infection: any sign of infection that will start 24 hours after surgery up

to 10 days. Include: post-surgical wound infection, endometritis and fever morbidity.

Endometritis: presence of fever accompanied by uterine tenderness, maternal tachycardia or

leukocytosis that happen postpartum.

Post-surgical Wound infection: infection on operation scar that occurs within 30 days after

the operation.

Fever morbidity: temperature more than 37.8°C after 24hours post operatively

Elective caesarean section: a planned caesarean section done when the mother is not in

labour.

Emergency caesarean section: caesarean section done when the mother is in labour and

there is deterioration to either maternal or foetal condition.

Neonatal sepsis: defined as sepsis within the first 28 days of life.

Early neonatal outcomes adverse outcome: admission to special care unit or a neonatal unit

for sepsis within the first 10 days of life.

X

ABSTRACT

Background: Prophylactic antibiotic administration is used to prevent post-surgical infection after caesarean section. Studies have suggested that the timing of prophylaxis plays an important role (Classen et al., 1992). There have been observed differences in outcome based on timing of administration. This is different depending on one group giving before skin incision another after skin incision. There is evidence that before skin incision has less incidence of post-operative infection without adverse neonatal outcome (Baaqeel & Baaqeel, 2013; Costantine et al., 2008). Over the years, the role of the anaesthesiologist in administering prophylactic antibiotics has come to prominence. There is increasing need for anaesthesia providers to understand the rationale of antimicrobial prophylaxis.

<u>Objective:</u> To compare the effect of administration of prophylactic antibiotic within one hour before skin incision and administration after skin incision on the incidence of post-operative infection in caesarean section patients in Mulago Hospital.

<u>Design:</u> A Single blinded randomised clinical trial done in Mulago Hospital on patients booked for caesarean section in labour ward theatre. The total sample size was 464.

Procedure: Patients were randomly allocated a group number which placed them on either arms of the study. They received the same prophylactic antibiotic according to their allotment that is either within one hour before skin incision or after skin incision as per current standard of practice. They were followed up for post-operative infection up to 10 days post operation. Data was entered with EpiData 3.1 and analyzed with STATA version 12.

Results: A significant difference of 19.2% P value 0.022 was seen when comparing incidence of overall post operative infection being less when prophylaxis is given before skin incision. Endometritis was found to be significantly reduced when giving prophylaxis before skin incision P value 0.036, whereas the difference in incidence of wound infection was not significant.

<u>Conclusion:</u> Giving prophylactic antibiotic before skin incision reduces incidence of post operative infection more especially endometritis.